


Light pollution threatens the Balearic shearwater

Petrel fledglings leave the nest after dark, but these marine birds' maiden flights towards the sea are hampered by city lights. Many collide or fall to the ground where they are in danger of being run over and vulnerable to predators. Now, a study has evaluated the impact of light pollution on three species of petrels on the Balearic Islands, including the Balearic shearwater, the most threatened sea bird in Europe, and concludes that between 30 and 47% of colonies are exposed to high levels of light pollution.

SINC

9/6/2015 08:30 CEST



The Balearic shearwater is the most threatened sea bird in Europe. / Beneharo Rodríguez

This is not a new phenomenon. The dangers that fledglings of certain species of seabirds face when they emancipate themselves from their parents by launching themselves towards the sea at night are caused by light pollution: collisions, being run over and greater vulnerability to predators when they fall to the ground.

The problem affects more than 40 species worldwide, some on the

endangered species list such as the Balearic shearwater or Newell's shearwater

Some archipelagos such as Hawaii, La Reunion, the Azores and the [Canary Islands](#) have borne witness to this threat for quite some time. In fact, the problem affects more than 40 species worldwide, some on the endangered species list such as the Balearic shearwater or Newell's shearwater (native to Hawaii).

"During some of the breeding seasons in all of these areas more than 2,000 birds are recorded to have been grounded in only 20 days (duration of the emancipation period)", Airam Rodríguez tells SINC. Rodríguez is a researcher at the department of Evolutionary Ecology at the Doñana Biological Station (CSIC) and lead author of a study analysing the data from three rescue centres on Mallorca, Menorca and Ibiza between 1999 and 2013.

The study, published in the *Journal of Ornithology*, evaluates the effect of outdoor lighting of city roads, cars, streets and sports facilities on three species of petrels which breed on the Balearic archipelago and are under threat: the Balearic shearwater (*Puffinus mauretanicus*), Scopoli's shearwater (*Calonectris diomedea*) and the European storm-petrel (*Hydrobates pelagicus*).



A Scopoli's shearwater affected by nocturnal city lights. / Beneharo Rodríguez

An ever-greater threat

The calculations (from 304 birds rescued at an average distance of 4,833 metres from their nearest colonies) demonstrate that at least 1% of the fledglings produced each year by the population are found on the ground due to light pollution. "This impact is very low compared to that reported for other species on other archipelagos," comments Rodríguez, although he highlights that the Balearic shearwater is the most affected of the three species.

"A swift rescue is crucial for the future survival of the grounded birds," says researcher Airam Rodríguez.

According to the researchers, the low percentage can be due to two factors: the distribution of the colonies (mainly located in areas close to the coast, cliffs or islets); and the absence of awareness-raising and rescue campaigns involving the public on the Balearic Islands. "In the Canary Islands, for example, these actions have been running since 1990 and in Hawaii since 1978," observes the scientist.

Menorca is the island where most birds are found on the ground (147), followed by Ibiza (92), Mallorca (61) and Formentera (4). The researchers evaluated their physical condition and drew one conclusion: as the emancipation period advances, their physical condition (body condition) decreases and the likelihood of fatalities increases.

"In reality, the swiftness with which the grounded birds are rescued is crucial for their future survival," confirms the expert. The study details that 26 of the 304 grounded birds died over 15-year period on the Balearic Islands as a result of light pollution.

Researchers also analysed the levels of light pollution around the breeding colonies using a nocturnal satellite image. "Between 30% and 47% of the colonies (depending on the species) are exposed to high levels of light pollution," points out Rodríguez.

Although the number of grounded birds does not seem alarming, the team

considers it a priority to make efforts to reduce levels of light pollution to the lowest possible. Bearing in mind that these are endangered species, the scientists recommend establishing rescue campaigns specifically for the emancipation periods, particularly in the case of the Balearic shearwater. "This would avoid an additional cause of mortality for these endangered seabirds," concludes Rodríguez.

Reference:

Airam Rodríguez et al. "Artificial lights and seabirds: is light pollution a threat for the threatened Balearic petrels?" *Journal of Ornithology* May 2015 DOI 10.1007/s10336-015-1232-3

Copyright: **Creative Commons**

TAGS

PARDELA BALEAR | AVES MARINAS | CONTAMINACIÓN LUMÍNICA | COLISIONES |
SUELO |

Creative Commons 4.0

You can copy, distribute and transform the contents of SINC. [Read the conditions of our license](#)